



SEPPIC

wesource

ACTIVE SCIENCE TO EMPOWER BEAUTY

JUVENESSENCE™

Youth in Full Power

Metabolism reactivation of aging cells

ALARIA ESCULENTA

The Artic Winged Kelp



ALARIA ESCULENTA: a brown seaweed that gets its name from its Wing-shaped Ribbed Thallus




- Alaria can grow up to 2 meters long & few dozen centimeters large.
- It lives in **extreme conditions** in agitated but transparent waters, where **sea pressure can exceed 6T/m²**.
- It is made up of a **network of elastic fibers** (proteins & phycopolysaccharides), which:
 - **preserves tissue integrity** against compression &
 - **ensures cohesion & firmness of all the cell architecture** (similar role that **proteic fibers & elastin** play in the Skin).
- It is **sourced in Europe (France & Ireland)**

About
The Artic
Winged Kelp

SKIN AGING: A COMPLEX MECHANISM

Involvement of nucleus membrane & some «cell actors»




SKIN PROTEINS SYNTHESIS ARE UNDER THE CONTROL OF DIFFERENT “CELL ACTORS”: THE NUCLEUS, THE RIBOSOME & THE PROTEASOMES.

- **Nucleus directs protein synthesis by synthesizing messenger RNA (mRNA)** according to instructions provided by the DNA.
- The **mRNA is then transported to the cytoplasm** via the nuclear membranes pores.
- Once an mRNA molecule reaches the cytoplasm, **ribosomes translate the mRNA's genetic message** into the primary structure of a specific polypeptide & finally in proteins.
- Proteasomes insure the **elimination of degraded proteins in amino acids** in the aging cells.
- These could be used again as **raw materials for the synthesis of new proteins**.

JUVENESSENCE™

Characteristics



INCI name: *Caprylic/Capric Triglyceride – Alaria esculenta extract*

Use level: 1%

Solubility: Liposoluble

Form: Liquid (pale yellow to dark green liquid)

Preservatives: Free

Claims: Metabolism reactivation of aging cells

Approvals:



RSPO
MASSBALANCE CERTIFIED



ECOCERT Cosmetics: raw material approved by ECOCERT GREENLIFE, in conformity with Ecocert natural and organic cosmetic standard. / **COSMOS Approved:** raw material approved by ECOCERT GREENLIFE, in conformity with Cosmos standard. / **NATRUE Approved:** raw material certified according to Natrue standard. / **MassBalance Certified:** supplied under MassBalance certification BVC-RSPO-1-1972708497. Contributes to the production of certified sustainable palm oil. / **HALAL Certified:** BiotechMarine site & products have been audited & certified by HCS (Halal Certification Services) ; officially recognized by MUI (Indonesia), JAKIM (Malaysia), MUIS (Singapore), CICOT, and other respected Halal authorities around the world.

JUVENESSENCE™

Agenda



Introduction:

ALARIA ESCULENTA

Skin aging: a complex mechanism

JUVENESSENCE™: Characteristics

- Reduction of accelerated aging protein: **Progerin** (▶)
- **Metabolism reactivation** of aging cells (▶)
- Regulation of **photo-aging mediators** (▶)
- **Anti-aging** efficiency: *In vivo* studies (▶)





JUVENESSENCE™

Reduction of
Accelerated Aging Protein:
Progerin

wesource by **SEPPIC**

JUVENESSENCE™

Reduction of the accelerated aging protein: PROGERIN




THE NUCLEUS: One of the first targets of aging

Structural integrity of the nucleus

- > This integrity plays an **essential role** in the adequate functioning of the cell, especially in:

- the **gene transcription**
- the **DNA replication**
- the **chromatin remodeling**



JUVENESSENCE™

Reduction of the accelerated aging protein: PROGERIN



PROGERIN,
a degraded form of the nuclear
architectural protein, Lamin A, is
involved in the premature

Publication by BiotechMarine

International Journal of Cosmetic Science, 2011, 1-5

International Journal of Cosmetic Science, 2011, 1-5

doi: 10.1111/j.1468-2494.2011.00656.x

Quantitative assessment of lactate and progerin production in normal human cutaneous cells during normal ageing: effect of an *Alaria esculenta* extract

C. Verdy*, J.-E. Branka* and N. Mekideche†


*Efficience, 178 rue de Brest, 35000 Rennes and †Biotechmarine, Zone Industrielle, BP 65, 22260 Pontrieux, France

Received 12 October 2010, Accepted 13 March 2011

Keywords: ageing, *Alaria esculenta*, algae extract, ELISA, lactate, progerin, young and aged fibroblasts

A- Normal fibroblast nuclei

B- Progeria fibroblast nuclei



In conclusion, we have demonstrated
for the first time that it is **possible to quantify
Progerin in human normal cutaneous cells.**

We have also shown that **in selected
experimental conditions, Progerin production
was linked to cell aging.**

JUVENESSENCE™

Reduction of the accelerated aging protein: PROGERIN




ASSESSMENT OF THE PROGERIN PRODUCTION in keratinocytes

Protocol

- **Human keratinocyte model, cultured in 3D**, from a *young subject* and a *mature subject* (from abdominal plasties, performed on women aged 35 to 66 years old)
- **Incubation of products** (3D models were incubated 96H)
- **Assessment of effects: Progerin assay (ELISA)**

Results



Explanation on the mechanism of action in Progerin regulation



Progerin is found in higher quantity in keratinocytes of older donors than those of young donors (**2.6 times higher**)

JUVENESSENCE™

Reduction of the accelerated aging protein: PROGERIN




CONSEQUENCES OF PROGERIN ACCUMULATION

At nucleus level:

- Nucleus membrane integrity loss
- Nucleus malformation
- Increase of DNA alteration
- DNA repair process alteration
- Genes transcription process alteration

At the organism level, Progerin increase affects:

- Skin proteins
- Dermis & epidermis tissues



JUVENESSENCE™

Reduction of the accelerated aging protein: PROGERIN



ASSESSMENT OF JUVENESSENCE™ ON THE PROGERIN PRODUCTION

Protocol

> **Human keratinocytes model**, cultured in 3D, from 'young' & 'mature' subjects
(from abdominal plasties, on women aged from 35 to 66 years old)

> **Reference product: the Insuline**
(increases the enzyme activity involved in the Progerin formation).

> **Incubation of products** (3D models were incubated 96H):
 in the absence of the product (control)
 in the presence of the reference product
 in the presence of JUVENESSENCE™ at different concentration rates

> **Assessment of effects:**
Progerin dosage (by Elisa kit) & **proteins dosage** (Bradford)


JUVENESSENCE™

Reduction of the accelerated aging protein: PROGERIN



ASSESSMENT OF JUVENESSENCE™ ON THE PROGERIN PRODUCTION

Results



At 1%, JUVENESSENCE™:

- Down-regulates the Progerin production in aging cells:

-78,7%

(66 years old donors)



-63.2%

(35 years old donors)

- Brings Back the Progerin levels at the rates observed on young cells (35 years old)





JUVENESSENCE™

Metabolism Reactivation
of Aging Cells

wesource by **SEPPIC**


JUVENESSENCE™

Metabolism reactivation of aging cells




All along the aging process, the ribosomal, mitochondrial & the proteasome activities are affected.

These decreases disturb the whole “skin cell machinery”, and more specifically the skin protein synthesis mechanism, as well as the detoxification of aging proteins



JUVENESSENCE™

Metabolism reactivation of aging cells



JUVENESSENCE™ 1%
reactivates the metabolism of
aging cells by stimulating:

- the detoxification (**+24%**),
- the mitochondrial activity (**ATP: +42%**),
- malate shuttles (**+31%**),
- cell viability (UVB): **+20%**,
- the ribosomal activity (**+29%**).





JUVENESSENCE™

Regulation of Photo-aging Mediators

wesource by **SEPPIC**

Skin,

in direct contact with the environment, undergoes aging as a consequence of U.V. exposure & environmental damages.

- **Sunlight** has a profound effect on the skin **causing premature skin aging**.
- **Chronological aging & photo-aging** share fundamental molecular pathways.
- **UV** stimulates and activates various cells & tissues to produce & release cytokines that may play a **significant role in the process of photo-aging**.
- Indeed, **photo-aging is regulated by a delicate balance** between the deleterious effects of cytokines leading to photo-aging & their beneficial effects ameliorating UV-caused damages.

JUVENESSENCE™

Regulation of photo-aging mediators for a soothing benefit



JUVENESSENCE™ 1%

plays an important role to **maintain homeostasis** in actinically damaged skin by **regulating the production of mediators**, and more specifically:

Pro-inflammatory mediators:

Interleukins 1, 6 & 8 (-12% to -21%)


TNF α (-14%)

IFN γ (-8%)

Anti-inflammatory mediators:

Interleukins 10

Interleukins 12p70





JUVENESSENCE™

In Vivo Studies

wesource by **SEPPIC**

JUVENESSENCE™

Anti-aging benefits - *In vivo* study 1/2



ASSESSMENT of the EFFICACY
of a cream containing **JUVENESSENCE™ 1%**,
applied **ON THE CROW'S FEET AREA**

In vivo protocol

- Study during 28 days, on 20 subjects with periorbital wrinkles
- Average age: 50 years old
- Cream applied twice a day
- Measurements with **Cutometer**

Subject at **D0**



Subject at **D28**



CUTOMETER STUDY

Results



Firmness: +25% vs D0



Elasticity: +20% vs D0



of the **number of wrinkles**
for **65%** of the volunteers



of the **depth of wrinkles**
for **80%** of the volunteers

JUVENESSENCE™

Anti-aging benefits- *In vivo* study 2/2



ASSESSMENT of the EFFICACY of a cream containing **JUVENESSENCE™ 1%**, applied **ON THE CHEEKS**

In vivo protocol

- Study during 28 days, on 20 subjects with a skin lacking firmness
- Average age: 50 years old
- Cream applied twice a day, under dermatological control
- **At D28, assessment of skin quality & firmness** (self-evaluation)

Results at D28

Dermatologist
Firmness: +10%



Self-evaluation
90% of the volunteers
noticed an **increase in**
Firmness , after 28 days



JUVENESSENCE™

Anti-aging benefits - SELF ASSESSMENT



% OF THE PANEL
PERCEIVING A **BENEFIT**,
after 28 days of treatment

Results at D28

In vivo: crow's feet

Number of
wrinkles:
65%

Depth of
wrinkles:
80%

Radiance:
70%

Hydration:
85%



In vivo: cheeks

Skin Tonicity:
85%


Firmness:
90%

Skin Grain
Improvement:
85%

Hydration:
70%

JUVENESSENCE™

Summary





JUVENESSENCE™

Annexes

ALARIA ESCULENTA, the Winged Kelp

Biogeography

Alaria esculenta is a kelp of Arctic/cold-temperature distribution, with its Southern distributional limit in the East-Atlantic at the coasts of Brittany and stretching northwards up to the coasts of Spitsbergen, as far as the winter sea ice. (Kraan et al, 2000). This species is common in the Faeroes, Greenland, Iceland (where it is eaten raw or cooked), and its absence in the southern North Sea and English Channel supports the hypothesis that the distribution of *Alaria esculenta* is temperature controlled. Indeed, its southernmost habitat in Europe is Brittany, France, in waters that do not exceed 16°C in temperature. (Munda, 1977)

This perennial seaweed, member of the Alariaceae family (Phylum Ocrophyta, Class Phaeophyceae), populates sublittoral zones, between 3-10 m depth on exposed rocky shores, while in very exposed sites, on vertical, very steep rocks it has been recorded at a 35 m depth.

Its growth is extremely fast, compared to other seaweeds, reaching up to 10 cm per day (Kraan and Guiry, 1998). *Alaria esculenta* can reach a length up to 6 m. Its short stipe gives rise to a long, thin, ribbon-like, wavy blade with a well-defined rib. The color is olive-green, and when touched, the blade feels satiny and flexible.

Life on the rocky shore

In some biotopes of the Arctic waters, *Alaria* forms large canopies, like other large laminarias do, often found in dense underwater forests. But this is usually not the case for *Alaria* in Brittany, where it is often accompanied by *Laminaria flexicaulis* and *Corallina officinalis* (Dizerbo, 1949). Here, it is exposed to violent surf and to the great pressure that waves create, of over 6t/m². More than any other seaweed, *Alaria* is conditioned to constant pounding by the waves. Holding on to the sloppy rocks, it can withstand the punishment that over 8000 waves dish out over the period of one day. On such a rocky shore, too dangerous to offer safe footing, only the seagulls enjoy the mad choreography that *Alaria*'s ribbons engage into, back and forth on the surface of the water, tugged and tossed in a rhythm borrowed from the sea.

It is precisely this fate teasing dance, this constant movement of the algal fronds that enables *Alaria* to enhance photosynthesis, by reducing self-shading and generating light flecks. Contrary to what one might expect, productivity and growth reveal themselves very high on wave-beaten shores. The action of drag forces stimulates carbon uptake and seaweed competitors may be removed by whiplash, while the wave action can also reduce herbivore grazing (Lobban, 1994).



Both *Alaria*'s chemistry and its morphology are adapted for living in the surf zone. The fertile pinnae in which the reproductive cells mature, are borne at the base of the frond, as this location is safer than the tips of the main blade. This is not the case for other rockweeds living higher on the shore and less subject to savage wave action, where the reproductive cells are formed at the tips of the fronds (Carson, 1998). Despite the apparent fragility of the translucent olive-green blade, its tissue, a web of elastic fibres, contains large amounts of alginic acid and its salts, which create a tensile strength and elasticity able to withstand the pulling and pounding of the waves. It was shown that the blades of wave-exposed laminarias have greater proportions of polyguluronate (strong, rigid gels) than those of wave-sheltered specimens (Venegas et al 1993, Lobban, 1994) and large proportions of cell wall polysaccharids and different forms of these, with their distinctive gelling properties are found in different regions of seaweeds (blade vs stipe)

ALARIA ESCULENTA, the Winged Kelp

The science drop

Alaria's biomass has been described as carbohydrate rich, with the main carbohydrate, alginate, accounting for 42% of the dry matter. (Schiener, 2013; Nwosu, 2010). In addition to structural carbohydrates, alginate and cellulose, laminarin, mannitol and fucoidan contents have also reported to comprise a significant proportion (Schiener, 2013).

The protein level is of about 11%, which is considerable for a brown seaweed, since, generally, brown seaweeds have lower protein levels than the red and green seaweeds.

High level of moist and ash (approx. 25 %) were measured during winter months in the same study. The ash content of Laminariales consist mainly of the ions- sodium, potassium, calcium, magnesium – with chloride and sulfate as the main counter-ions. Along with these macro-nutrients, trace metals such as Fe, Zn, Mn, Al and Cu have been identified. (Schiener, 2013; Schumacher, 2011)

Alaria esculenta was shown to be a species of high phenotypic plasticity, its acclimating capacities enabling it to respond to short-term stresses. As Wienke and Bischof show (Seaweed Biology, 2012), seaweeds populating a flexible environment require mechanisms of acclimation in order to set physiological performance to the variation of environmental requirements. Acclimation to UV exposure has been demonstrated in Alaria esculenta. Under repeated UV exposure, the competence of recovery from UV-induced photoinhibition increased after just a few exposure/recovery cycles, which might indicate an activation of different repair mechanisms, counteracting the impact of UV exposure by a faster replacement of damaged molecules. Moreover, the degree of inhibition became smaller, which might also be related to an activated ROS defense system counteracting UVB mediated oxidative stress, or the formation of UV-screening compounds, like phlorotannins. Indeed, an increase in the size of phlorotannin containing vesicles (physodes) was observed in Alaria esculenta after UVB exposure, indicating an induction of phlorotannin synthesis (C. Wienke and K. Bischof, 2012)

Seaweeds, as photosynthetic organisms, are exposed to a combination of light and high oxygen concentration at the origin of the formation of free radicals and other oxidative reagents. But the awareness of the lack of structural damage in their organs has led the scientific community to consider that their protection against oxidation comes from their natural content, or production under stress, in antioxidant substances. Indeed, macroalgae are particularly rich in natural antioxidants, e.g. phlorotannins, ascorbic acid, tocopherols and carotenoids. In a study

carried on several Phaeophyta from the Brittany coast (Zubia et al, 2009) a significant correlation was found between the high antioxidant activity and the high phenolic content. Alaria esculenta was shown in this study to possess important antioxidant and antitumoral activities. Further anti-proliferative and anti-diabetic effects were discussed in a study by Nwosu et al, 2010. The multifunctional antioxidant activity of polyphenols was discussed by Wang et al 2008 in a study that invites to broadening the reflexion towards the research of synergistic effects of other active components than just phlorotannins, like maybe low-molecular-weight polysaccharides, proteins or peptides that could also contribute to the scavenging effect.

As determined for the Arctic Kongsfjord (Lippert et al 2001), Alaria esculenta is a species of utmost ecological relevance, hosting up to 51 epifauna species. Here, they act as food, habitat and nursery for multiple associated organisms (Fredersdorf et al 2009). On the Brittany coasts, the ecology of Alaria esculenta has not yet been precisely documented, but a close look at its growth sites would marvel the eye, for if at high tide nothing but Alaria's fronds show tossing in the wind, at low tide, when the waters retrieve, a kaleidoscope of colors and forms show their beauty along the rocks. The pink corallina algae, the electric blue Irish moss, the yellow sponge, brown rockweeds, or sea anemones like the solitary and very fashionable Sagartia elegans, all delicately stir with life, in a world of crushing force.

By Anca BOATA

ALARIA ESCULENTA, the Winged Kelp

Bibliography


1. R. Carson, *The Edge of the Sea*, Mariner Books 1998.
2. H. Dizerbo, *Algues Marines. Quelques aspects de la vegetation des Algues marines supérieures du Massif Armoricaïn*, 1949
3. J. Fredersdorf et al, *Interactive effects of radiation, temperature and salinity on different life history stages of the Arctic kelp Alaria esculenta*, 2009
4. C.S. Lobban and PJ Harisson, *Seaweed Ecology and Physiology*, 1994
5. M. Jaspars and F. Folmer, *Sea vegetables for Health*, 2013
6. S. Kraan and M.D. Guiry, *Strain selection in the edible brown seaweed Alaria esculenta: Genetic fingerprinting and hybridization studies under laboratory conditions*, 1998
7. S. Kraan et al, *The edible brown seaweed Alaria esculenta (Phaeophyceae, Laminariales): hybridization, growth and genetic comparaisn of six Irish populations*, 2000
8. H. Lippert et al, *Macrofauna associated with macroalgae in the Kongsfjord (Spitsbergen)*, 2001
9. K. Mai et al, *Comparative studies on the nutrition of two species of abalone. II. Amino acid composition of abalone and six species of macroalgae with an assessment of their nutritional value*, 1994
10. I. Munda and K. Luning, *Growth performance of Alaria esculenta off Helgoland*, 1977
11. F. Nwosu et al, *Anti-proliferative and potential anti-diabetic effects of phenolic-rich extracts from edible marine algae*, 2010
12. P. Schiener et al, *The seasonal variation in the chemical composition of the kelp species Laminaria digitata, Laminaria Hyperborea, Saccharina latissima and Alaria esculenta*, 2014
13. M. Schumacher et al, *Hydrothermal conversion of seaweeds in a batch autoclave*, 2011
14. T. Wang et al, *Total phenolic compounds, radical scavenging and metal chelation of extracts from Icelandic seaweeds*, 2009
15. C. Wiencke and K. Bischof, *Seaweed Biology. Novel insights into Ecophysiology, Ecology and Utilization*, 2012
16. Q. Zhang et al, *A simple 96-well microplate method for estimation of total polyphenol content in seaweeds*, 2006
17. M. Zubia et al, *Antioxidant and antitumoral activities of some Phaeophyta from Brittany coasts*, 2009
18. <http://www.jncc.gov.uk/marine/biotopes> Joint Nature Conservation Committee website, On the biotope of Alaria esculenta and the associated epifauna



JUVENESSENCE™


SENESCENCE AMPLIFIER & MTOR

CELL membrane



Overactivation of mTOR pathway causes aging disorders.

With age and stress, **mTOR production increases**, inhibiting the autophagy, the elimination of damaged proteins, as Progerin.



mTOR protein

(Target Of Rapamycin mTOR): intracellular enzyme, discovered in 1994, having a kinase activity.

Essential role in the development & aging processes

Cellular Senescence under the mTOR control.

A reduction of the activity of the **mTOR signaling pathway** increases the lifespan of various organisms

Activation by a membrane receptor, activated in its turn by signals (nutriments, stress...)

JUVENESSENCE™

ACTIVITY OF JUVENESSENCE™ ON mTOR




Protocol

- Monolayer of human dermal fibroblasts: biopsies from women aged 29 & 47 years old
- Reference inhibitor: Rapamycin 100 nM (helps Progerin elimination from cells)
- Cells are incubated 96h, without the reference product, with the reference product, and with JUVENESSENCE™ (1% v/v)
- Dosage of mTOR: quantification in cell medium by Elisa (specific)

Results

Comparison of mTOR production between young & mature human fibroblasts in culture



mTOR production is significantly superior in mature human cells than in young ones.


JUVENESSENCE™

ACTIVITY OF JUVENESSENCE™ ON mTOR



Results

Action of **JUVENESSENCE™ 1%** on mTOR production in mature human fibroblasts



JUVENESSENCE™ 1%

- inhibits mTOR production: **-68,5%**

Reference Rapamycin

- inhibits mTOR production: **-74%**

JUVENESSENCE™ 1% decreases significantly the **mTOR production** in mature human cells.

Return
slide 9



BIBLIOGRAPHY



“Rapamycin activates autophagy in Hutchinson-Gilford progeria Syndrome. Implications for normal aging and age-dependent neurodegenerative disorders”

Autophagy 8:1, 147–151; January 2012; G 2012 Landes Bioscience.

John J. Graziotto,¹ Kan Cao,² Francis S. Collins³ and Dimitri Krainc¹

¹ Department of Neurology; Massachusetts General Hospital; MassGeneral Institute for Neurodegenerative Disease; Harvard Medical School; Charlestown, MA USA;

² Department of Cell Biology and Molecular Genetics; University of Maryland; College Park, MD USA;

³ Genome Technology Branch; National Human Genome Research Institute; National Institutes of Health; Bethesda, MD USA

“Rapamycin Reverses Cellular Phenotypes and Enhances Mutant Protein Clearance in Hutchinson-Gilford Progeria Syndrome Cells”

Research article, 29 June 2011 Vol 3 Issue 89 89ra58 / www.ScienceTranslationalMedicine.org

Kan Cao^{1,2}, John J. Graziotto³, Cecilia D. Blair¹, Joseph R. Mazzulli³, Michael R. Erdos¹, Dimitri Krainc³, Francis S. Collins¹

¹Genome Technology Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD 20892–8004, USA.

²Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD 20742, USA.

³Department of Neurology, Massachusetts General Hospital, MassGeneral Institute for Neurodegenerative Disease, Harvard Medical School, Charlestown, MA 02129, USA.

“Lamin A-dependant misregulation of adult stem cells associated with accelerated ageing”

PubMed Central Journal List; Nat Cell Biol. 2008 April; 10(4): 452-459

Paola Scaffidi¹ and Tom Misteli¹

¹ National Cancer Institute, NIH, Bethesda, Maryland 20892, USA

“Lamin A mutation and the cell cycle, a connection between Progeria and normal aging”

www.ouroboros.wordpress.com/2007/04/03

“Progerin and telomere dysfunction collaborate to trigger cellular senescence in human fibroblasts”

www.jci.org/articles/view/43578

J. Clin Invest. 2011; 121(7):2833-2844. Volume 121, issue 7 (July 1, 2011) / American Society for Clinical Investigation^{1,2}

Cecilia D. Blair¹, Dina A. Faddah¹, Julia E. Kieckhafer², Michelle Olive¹, Michael R. Erdos¹, Elizabeth G. Nabel¹ and Francis S. Collins¹

¹ Genome Technology Branch, National Human Genome Research Institute, NIH, Bethesda, Maryland, USA.

² Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, Maryland, USA.



SEPPIC

wesource

ACTIVE SCIENCE TO EMPOWER BEAUTY

**Thank you
for your attention**

wesourcebeauty.com

A company
Air Liquide
HEALTHCARE

Contacts



Nota

The analytical specifications warranted are only those mentioned on the certificate of analysis supplied with each delivery of the product.

Except as set forth above, SEPPIC* makes no warranties, whether express, implied or statutory, as to the product which is the subject of this document. Without limiting the generality of the foregoing, SEPPIC* makes no warranty of merchantability of the product or of the fitness of the product for any particular purpose. Buyer assumes all risk and liability resulting from the use or sale of the product, whether singly or in combination with other goods. The information set forth herein is furnished free of charge and is based on technical data that SEPPIC* believes to be reliable. It is intended for use by persons having technical skill and at their own discretion and risk. Since conditions of use are outside SEPPIC*'s control, SEPPIC* makes no warranties, express or implied, and assumes no liability in connection with any use of this information. Nothing herein is to be taken as a license to operate under or a recommendation to infringe any patents.

* SEPPIC being:

SEPPIC S.A. - Paris establishment

Paris La Défense
50 boulevard National
CS 90020
92257 La Garenne Colombes Cedex
FRANCE
Tél : +33 (0)1 42 91 40 00
info.seppic@airliquide.com

SEPPIC S.A. - Castres establishment

127, chemin de la poudrerie – BP 90128
81105 Castres Cedex
FRANCE
Tél : +33 (0)5 63 72 69 69
Fax : +33 (0)5 63 72 69 70

BiotechMarine S.A.S.

Zone Industrielle
B.P. 72
22260 Pontrieux
FRANCE
Tél : +33 (0)2 96 95 31 32
contact.biotechmarine@airliquide.com

Serdex S.A.S. - Pau establishment

1 avenue Normandie Niemen
64140 LONS
FRANCE
Tél : + 33 (0)5 59 13 04 20
contact-serdex@airliquide.com

UNITED STATES - SEPPIC Inc.

30, Two Bridges Road, suite 210
Fairfield
New Jersey 07004-1530
UNITED STATES
Tél : +1 973 882 5597
us.seppic@airliquide.com

UNITED STATES - Polykon Manufacturing LLC

6201 Engineered Wood Way
Sandston
Henrico County VA 23150
UNITED STATES
Tél : +1-804-624-9057
sharon.harper@airliquide.com

BRAZIL - SEPPIC Brasil EQCI Ltda

Alameda Santos 745/ cj 72/ 7th floor
Cerqueira Cesar
01419-001
São Paulo – SP
BRAZIL
Tél : +55 11 3252 3911
brasil.seppic@airliquide.com

COLOMBIA - SEPPIC Colombia SAS

Calle 71 n°10-40
Edificio Orbe 71 of 401
Bogota
COLOMBIA
Tél : +571 702 44 48
colombia.seppic@airliquide.com

GERMANY - SEPPIC GmbH

Martinstrasse 16-20
50667 Köln
GERMANY
Tél : +49 (0)221 888824 0
germany.seppic@airliquide.com

ITALY - SEPPIC Italia Srl

Via Quarenghi 27
20151 Milano
ITALY
Tél : +39 02 38009110
italy.seppic@airliquide.com

POLAND - SEPPIC Office

Schülke Polska Sp. z o.o.
Al. Jerozolimskie 132
02-305 Warsaw
POLAND
Tel. +48 22 11 60 759
poland.seppic@airliquide.com

RUSSIA - SEPPIC office

OOO AIR LIQUIDE
Vorontsovskaya St. 17
109147 Moscow
RUSSIA
Tel. +7 495 641 28 98 ext. 375

CHINA - SEPPIC Shanghai Chemical Specialities Co.

819 West Nan Jing Road
Shanghai 200041
CHINA
Tél : +86 (21) 64 66 01 49
china.seppic@airliquide.com

INDIA - Air Liquide India Holding Pvt Ltd

O2 Commercial Twin Tower
B/1304, Asha Nagar Road
Mulund (W)
Mumbai 400080
INDIA
Tél : + 91 22 61046464
india.seppic@airliquide.com

SINGAPORE - SEPPIC Office

Jem Office Tower
52 Jurong Gateway Road
#15-02 - Singapore 608550
SINGAPORE
Tél : +65 6373 7221
singapore.seppic@airliquide.com

JAPAN - SEPPIC Office

Air Liquide Japan's Office
Granpark Tower 3-4-1 Shibaura
Minato-ku, Tokyo 108-8509
JAPAN
Tél : +81 3 6414 6725
japan.seppic@airliquide.com

KOREA - SEPPIC Office

12FL, V-Plex,
501, Teheran-ro
Gangnam-gu,
Seoul, SOUTH KOREA 06168
Tél : +82-2-3019-2569
korea.seppic@airliquide.com

DUBAI - SEPPIC Office

East Wing 5,
Block-A, 3rd Floor
Dubai Airport Free Zone
UNITED ARAB EMIRATES
Tél : +971 (4) 20 55 500
dubai.seppic@airliquide.com